

## 17. Soft tissue sarcomas


Total number of cases in the registry (1980-2004): 2454

5-, 10-, and 15-year survival probabilities (life table estimates according to (8)): 64 %, 60 %, and 57 %


### Selected characteristics (Germany: 2000-2004)

|                                             |                    |               |
|---------------------------------------------|--------------------|---------------|
| Relative frequency:                         | 615 / 8948 = 6.9 % |               |
| Incidence rates (per 100,000):              | age-standardized   | cumulative    |
| girls (n = 280)                             | 0.9                | 14.0          |
| boys (n = 335)                              | 1.1                | 16.1          |
| total (n = 615)                             | 1.0                | 15.1          |
| Sex ratio (boys to girls):                  | 1.2 : 1            |               |
| Age-specific incidence rates (per 100,000): |                    |               |
| less than 1 year (n = 78)                   | 2.1                |               |
| 1 - 4 years (n = 202)                       | 1.3                |               |
| 5 - 9 years (n = 142)                       | 0.7                |               |
| 10 - 14 years (n = 193)                     | 0.8                |               |
| Age:                                        | 25%-quantile       | 2 5/12 years  |
|                                             | median             | 5 8/12 years  |
|                                             | 75%-quantile       | 11 2/12 years |
| Relative frequency of trial patients:       | 93 %               |               |

### Age- and sex-specific incidence rates (Germany: 2000-2004)



### Age-standardized annual incidence rates (Germany: 1980-2004)



### Survival probabilities for different soft tissue sarcomas (Germany: 1980-2004)

